

In the Claims:

Amend Claims 82, 83, 87, 88, 91, 94, and 97-100 as indicated below.

82. (Amended) A method for stimulating an immune response in an animal, the immune response being directed toward a self-epitope [antigen], the method comprising:

- Sub E1
D1
- a) providing a ubiquitin fusion protein comprising ubiquitin fused to one or more [contiguous or non-contiguous] epitope-containing segments at contiguous or non-contiguous locations within ubiquitin, the epitope-containing segments comprising two or more identical or non-identical non-ubiquitin self-epitopes, wherein the ubiquitin fusion protein is immunogenic for the non-ubiquitin self-epitopes therein; and
 - b) administering the ubiquitin fusion protein of step a) to an animal under conditions appropriate for the stimulation of an immune response, thereby stimulating an immune response to the non-ubiquitin self-epitopes.

83. (Amended) The method of Claim 82 wherein the ubiquitin fusion protein of step a) is further conjugated to an immunogenic carrier protein prior to administration.

Sub E2
D2

87. (Amended) The method of Claim 85 wherein the ubiquitin [moiety] is modified such that the epitope-containing segment fused to the C-terminus of ubiquitin is non-cleavable by a ubiquitin-specific protease.

88. (Amended) The method of Claim 84 wherein the non-ubiquitin self-[antigen]epitope is a peptide hormone.

Sub E3
D3

91. (Amended) The method of Claim 90 wherein the physiological consequence[s] of administration of the ubiquitin fusion protein to the animal is immunocastration of the animal [are

sub E3 D3
cut int
substantially similar to the consequences of surgical castration].

94. (Amended) The method of Claim 90 wherein two [non-contiguous] epitope-containing segments, each comprising the amino acid sequence QHWSYGLRPGQHWSYGLRPG (SEQ ID NO: 26) are fused to a single ubiquitin molecule at non-contiguous locations within ubiquitin, one being fused to the N-terminal amino acid of ubiquitin, the other being fused at an internal site of ubiquitin, the internal site being between residue 35 and 36 of ubiquitin.

97. (Amended) A method for stimulating an immune response in an animal, the immune response being directed toward a [ubiquitin fusion protein which contains a] self-[antigen] epitope, the method comprising:

- sub E5
D5
- a) providing a ubiquitin fusion protein comprising ubiquitin fused to a single epitope-containing segment, the epitope-containing segment comprising two or more identical non-ubiquitin self-epitopes, wherein the ubiquitin fusion protein is immunogenic for the non-ubiquitin self-epitopes therein;
 - b) administering the ubiquitin fusion protein of step a) to an animal under conditions appropriate for the stimulation of an immune response, thereby stimulating an immune response to the non-ubiquitin self-epitopes.

98. (Amended) A method for stimulating an immune response in an animal, the immune response being directed toward a [ubiquitin fusion protein which contains a] self-[antigen]epitope, the method comprising:

- a) providing a ubiquitin fusion protein comprising ubiquitin fused to two or more [non-contiguous] epitope-containing segments, at contiguous or non-contiguous locations within ubiquitin, each epitope-

containing segment comprising one or more identical or non-identical non-ubiquitin self-epitopes, wherein the ubiquitin fusion protein is immunogenic for the non-ubiquitin self-epitopes therein;

- b) administering the fusion protein of step a) to an animal under conditions appropriate for the stimulation of an immune response, thereby stimulating an immune response to the non-ubiquitin self-epitopes.

99. (Amended) A method for stimulating an immune response in an animal, the immune response being directed toward a [ubiquitin fusion protein which contains a] self-[antigen]epitope, the method comprising:

- a) providing a ubiquitin fusion protein comprising ubiquitin fused to a single epitope-containing segment comprising two or more identical or non-identical non-ubiquitin self-epitopes, the epitope-containing segments being fused to ubiquitin at fusion sites selected from the group consisting of the N-terminus and an internal fusion site, wherein the ubiquitin fusion protein is immunogenic for the non-ubiquitin self-epitopes therein;
- b) administering the ubiquitin fusion protein of step a) to an animal under conditions appropriate for the stimulation of an immune response, thereby stimulating an immune response to the non-ubiquitin self-epitopes.

100. (Amended) A method for stimulating an immune response in an animal, the immune response being directed toward a [ubiquitin fusion protein which contains a] self-[antigen]epitope, the method comprising:

- a) providing a ubiquitin fusion protein comprising ubiquitin fused to a single epitope-containing segment comprising one or more identical or non-identical non-ubiquitin self-epitopes, the epitope-containing segment

mb
CS
wk

DS
wk